

FOLIAR APPLICATION OF GLYCINE BETAINE AMELIORATED DROUGHT STRESS AND ENHANCED THE BIOMASS AND OTHER GROWTH-LINKED BIOCHEMICAL AND PHYSIOLOGICAL ATTRIBUTES IN WHEAT

HAFIZ MUHAMMAD WAQAS¹, MUHAMMAD AKBAR^{1*}

¹*Department of Botany, University of Gujrat, Punjab, 50700, Pakistan.*

**Corresponding Author's email: muhammad.akbar@uog.edu.pk*

Abstract:

Drought stress causes significant losses in crop yield and other parameters of wheat crops. Glycine betaine (GB) is found in plants naturally and helps the plant to cope with abiotic stress. In the present investigation effect of foliar application of Glycine betaine under drought stress was evaluated on wheat plants by adopting a completely randomized design (CRD). The effect of six treatments viz Control, 80% field moisture capacity (FMC), 60% FMC, 5 mM GB, 80% FMC+5mM GB, and 60% FMC+5mM GB were investigated on morphological, biochemical, and physiological attributes of wheat. Both 80 and 60% FMC showed a significant reduction in all these parameters but this decline is more pronounced at the former level of drought. The foliar application of GB significantly enhanced the shoot length, shoot fresh weight, chlorophyll, carotenoids, photosynthetic rate (*Pn*), transpiration rate (*E*), and stomatal conductance (*gs*) by 24, 26, 10, 12, 16, 20, and 16% respectively as compared to control. Combined application of GB with drought stress also enhanced the plant height and shoot fresh weight by 28 and 41%, respectively, Chl a, b and carotenoids by 10, 10, and 12%, *Pn*, *E*, and *gs* by 27, 47, and 35%, respectively as compared to 60% FMC drought treatment level. So based on these results, it is concluded that GB is an effective organic osmolyte that have promising potential to ameliorate drought stress and enhanced the biomass and other growth-linked biochemical and physiological attributes.

Keywords: Drought stress, Morphology, Photosynthetic, Stomatal conductance.

Introduction

A major threat to ecosystems, agriculture, water resources, and human well-being worldwide is drought stress, a ubiquitous and growing environmental concern (Sakamoto & Murata, 2002; Zia *et al.*, 2021). Drought stress is characterized by extended periods of exceptionally low precipitation and water availability, and it can have significant effects on many facets of life on Earth (Abbass *et al.*, 2022). The frequency, intensity, and duration of drought events are growing as global climate patterns continue to change, amplifying their negative consequences on both natural and human systems (Weiskopf *et al.*, 2020).

Drought stress can reduce crop yields, delay plant growth, and even cause crop failure in agricultural systems (Hussain *et al.*, 2018). These errors interrupt food production, increase food scarcity, and boost food prices, resulting in an increase in food poverty around the world. Drought stress greatly affects the amount of available water by reducing surface water

bodies, groundwater reserves, and soil moisture (Ahluwalia *et al.*, 2021). This scarcity has an influence on both industrial operations and human consumption, resulting in water rationing, water-borne infections, and severe economic repercussions in a variety of businesses (Saha *et al.*, 2022). Drought stress destabilizes ecosystems by altering plant diversity, reducing biodiversity, and compromising the health of wetlands, forests, and other natural habitats. These alterations have a negative impact on key functions (Hussain *et al.*, 2018; Ahmad *et al.*, 2022).

The world's food supply is threatened by declining water resources and challenged by rising dietary needs (Hanjra & Qureshi, 2010). The primary economic driver in Pakistan is agriculture, and among the major crops, rice dominates. Agriculture is the economic backbone of Pakistan. Pakistan's basic food is wheat, and as the country's population grows, so does the demand for food. A significant abiotic stress that can negatively affect wheat productivity is drought (Rehman *et al.*, 2015). Wheat plants are

susceptible to it at every stage of development, from germination to maturity (EL Sabagh *et al.*, 2021). The length and intensity of the drought, as well as the genetics of the wheat cultivar, will all have an impact on how severe the effect is (Tricker *et al.*, 2018; Lama *et al.*, 2023).

Glycine betaine (GB) is a naturally occurring chemical molecule found in a variety of plants (Ali *et al.*, 2020). It is an osmolyte, which implies that it protects cells from dehydration. Glycine betaine can also help plants resist additional stressors like heat, cold, and salinity (Sakamoto & Murata, 2002). Previous studies reported that GB can trigger the osmotic potential in plants by regulating the reactive oxygen species (ROS). Previous studies revealed that GB can ameliorate the drought effect on different crops but very few studies were present regarding wheat. So, the objective of this study is to check the drought mitigation potential of GB on wheat plants attributes.

Material and methods

To investigate the effect of drought stress and foliar application of Glycine betaine (GB) on wheat crops an experiment was conducted in Gujarat, Punjab, Pakistan. Wheat verity 2008 was grown in plastic pots having a diameter of 20 cm and a height of 25 cm. The pots were filled with 8 kg of soil. % seeds were sown in each pot and on germination 2 plants were kept to check the effect of GB and drought.

Treatments: Following treatments were used to assess the effect of GB and drought stress on wheat plants.

T1= Control (100% Field Moisture Capacity (FMC)

T2= Wheat + 80% FMC

T3= Wheat+ 60 % FMC

T4= Wheat+ 5 mM Glycine betaine (GB)

T5= Wheat+ 80% FMC+5 mM GB

T6= Wheat+ 60% FMC+5 mM GB

Each pot was sprayed with 10 ml of 5mMGB spray. Pots were kept under greenhouse conditions to avoid rain and maintain the drought level.

Morphological Parameters: Morphological parameters were assessed after 25 days of treatment application. Shoot length root length and their fresh and dry weight was measured by physical means and electrical balance.

Biochemical Parameters: To determine the chlorophyll a, b, and carotenoids similar method was adopted as described by Arnone (1949) and Devis (1976).

Physiological Parameters: In physiological parameters, net photosynthetic rate (Pn), stomatal conductance (gs), and transpiration rate were measured with the help of an Infrared gas analyzer machine on full sunny days between 11:00 am to 2:00 pm.

Results and Discussion

Effect of Drought stress and GB on morphology of wheat: Drought stress significantly reduced the plant height of wheat plants by 27 and 48% respectively as compared to healthy control. On the other hand, foliar application of Glycine betaine (GB) increased the plant height by 24% as compared to the untreated healthy control. Moreover, foliar application of GB applied in combination with drought stress plant height significantly increased by 11 and 28% respectively as compared to drought-bearing plants (Figure 1A).

Drought stress significantly reduced the shoot fresh weight of wheat plants at both drought levels 80 and 60% FMC. However, this decline is more prominent at the 60% FMC level. On the other hand, the Foliar application significantly enhanced the fresh weight by 26% when applied alone. Furthermore, the combined application of GB with 80% FMC increased the fresh weight of wheat plants by 25%, and with 60% FMC, a prominent increase of 41% was recorded as compared to their alone drought level (Figure 1 B).

Plant height reduces significantly with an increase in the severity of drought conditions. This reduction in the height may be a result of inhibition of cell elongation by severe drought stress (Yang *et al.*, 2021). Drought causes impaired germination and poor establishment of plants. Cell growth is one of the most important process that is sensitive to drought. It reduces due to decreases in turgor pressure. Growth results from the production of daughter cells from meristematic cell division and the expansion of young cells (Osmolovskaya *et al.*, 2018). Under severe conditions of drought, cell division can be inhibited

by interruption of water flow from the xylem to elongating cells, growth, and yield (Wahab *et al.*, 2022). A very common and damaging effect of drought stress on crop plants is the reduction in fresh and dry weight (Zia *et al.*, 2021). Drought causes a decrease in the number of leaves per plant, individual leaf size, and longevity of leaves. Expansion in the leaf area depends upon the turgor pressure, temperature, and sufficient supply of nutrients for growth. Reduced leaf area under water stress is damaging because it leads to a reduction in nutrient uptake due to reduced transpiration (Fahad *et al.*, 2017).

Effect of Drought and Glycine Betain on wheat biochemicals and physiological attributes :Drought stress depicted the negative effect on photosynthetic pigments chlorophyll a, b and carotenoid and reduced them by 57, 59 and 67%, respectively, when plants were grown at 60% FMC as compared to the control. On the other hands, GB showed statistically significant results and showed increased trend in Chl a, b, and carotenoid pigments. While drought and GB combined affect also showed significant increase of 40% in Chl a, a 43% uplift in Chl b and 53% increased in carotenoid content (Figure 2 A-C).

In the present investigation drought stress significantly affect the gaseous exchange parameters and significantly reduced the photosynthetic rate (Pn) by 57 %, transpiration rate (E) by 70% and stomatal conductance (gs) by 56% as compared to control. Foliar application of GB can enhance the Pn and gs by 16% and E by 20%, respectively, as compared to healthy control. On the other hand, combined application significantly mitigated the drought stress and enhanced the Pn, E, and gs by 27, 47 and 56% against different FMC drought levels (Figure 3 A-C)

Drought stress reduces the pigments like chlorophyll a and chlorophyll b. Drought stress created changes in the fraction of chlorophyll 'a' and 'b' and carotenoids (Shanthi *et al.*, 2023). A decrease in chlorophyll content was testified in drought-stressed cotton. Drought stress decreases more chlorophyll content than chlorophyll b and total carotenoids (Zhuang *et al.*, 2020). As water is most important element of photosynthetic activity and limiting of water impair the photosynthetic rate. Drought stress greatly decreased stomatal conductance (gs), and transpiration rate (E). And due to the decrease of these activities, photosynthesis also decreases (Seleiman *et al.*, 2021). Plants retort to stresses such as drought by decreasing leaf expansion

and closing stomatal pores. In this way, they save the water, nutrients, and carbohydrates necessary for survival, which is helpful for sustaining optimal water status, photosynthesis, and plant growth (Dos Santos *et al.*, 2022).

References

Abbass, K., Qasim, M. Z., Song, H., Murshed, M., Mahmood, H., & Younis, I. (2022). A review of the global climate change impacts, adaptation, and sustainable mitigation measures. *Environmental Science and Pollution Research*, 29(28), 42539–42559. <https://doi.org/10.1007/s11356-022-19718-6>

Ahluwalia, O., Singh, P. C., & Bhatia, R. (2021). A review on drought stress in plants: Implications, mitigation and the role of plant growth promoting rhizobacteria. *Resources, Environment and Sustainability*, 5, 100032. <https://doi.org/10.1016/j.resenv.2021.100032>

Ahmad, S., Zahra, J., Ali, M., Ali, S., Iqbal, S., Kamal, S., Tahir, M., & Aborode, A. T. (2022). Impact of water insecurity amidst endemic and pandemic in Pakistan: Two tales unsolved. *Annals of Medicine & Surgery*, 81. <https://doi.org/10.1016/j.amsu.2022.104350>

Ahmed, N., Zhu, M., Li, Q., Wang, X., Wan, J., & Zhang, Y. (2021). Glycine betaine-mediated root priming improves water stress tolerance in wheat (*Triticum aestivum* L.). *Agriculture*, 11(11): 1127. <https://doi.org/10.3390/agriculture1111127>

Ali, S., Abbas, Z., Seleiman, M. F., Rizwan, M., Yavaş, İ., Alhammad, B. A., Shami, A., Hasanuzzaman, M., & Kalderis, D. (2020). Glycine betaine accumulation, significance and interests for heavy metal tolerance in plants. *Plants*, 9(7), 896. <https://doi.org/10.3390/plants9070896>

Dos Santos, T. B., Ribas, A. F., De Souza, S. G. H., Budzinski, I. G. F., & Domingues, D. S. (2022). Physiological responses to drought, salinity, and heat stress in plants: A review. *Stresses*, 2(1), 113–135. <https://doi.org/10.3390/stresses2010009>

EL Sabagh, A., Islam, M. S., Skalicky, M., Ali Raza, M., Singh, K., Anwar Hossain, M., Hossain, A., Mahboob, W., Iqbal, M. A., Ratnasekera, D., Singhal, R. K., Ahmed, S., Kumari, A., Wasaya, A., Sytar, O., Brestic, M., ÇIG, F., Erman, M., Habib Ur Rahman, M., ... Arshad, A. (2021). Salinity stress in wheat (*Triticum aestivum* L.) in the changing climate: Adaptation and management strategies. *Frontiers in Agronomy*, 3. <https://www.frontiersin.org/articles/10.3389/fagro.2021.661932>

Fahad, S., Bajwa, A. A., Nazir, U., Anjum, S. A., Farooq, A., Zohaib, A., Sadia, S., Nasim, W., Adkins, S., Saud, S., Ihsan, M. Z., Alharby, H., Wu, C., Wang, D., & Huang, J. (2017). Crop production under drought and heat stress: Plant responses and management options. *Frontiers in Plant Science*, 8. <https://www.frontiersin.org/articles/10.3389/fpls.2017.01147>

Hanjra, M. A., & Qureshi, M. E. (2010). Global water crisis and future food security in an era of climate change.

Food Policy, 35(5), 365–377. <https://doi.org/10.1016/j.foodpol.2010.05.006>

He, C., Zhang, W., Gao, Q., Yang, A., Hu, X., & Zhang, J. (2011). Enhancement of drought resistance and biomass by increasing the amount of glycine betaine in wheat seedlings. *Euphytica*, 177(2), 151–167. <https://doi.org/10.1007/s10681-010-0263-3>

Hussain, H. A., Hussain, S., Khaliq, A., Ashraf, U., Anjum, S. A., Men, S., & Wang, L. (2018). Chilling and drought stresses in crop plants: Implications, cross talk, and potential management opportunities. *Frontiers in Plant Science*, 9, 393. <https://doi.org/10.3389/fpls.2018.00393>

Lama, S., Leiva, F., Vallenback, P., Chawade, A., & Kuktaite, R. (2023). Impacts of heat, drought, and combined heat–drought stress on yield, phenotypic traits, and gluten protein traits: Capturing stability of spring wheat in excessive environments. *Frontiers in Plant Science*, 14. <https://www.frontiersin.org/articles/10.3389/fpls.2023.1179701>

Osmolovskaya, N., Shumilina, J., Kim, A., Didio, A., Grishina, T., Bilova, T., Keltsieva, O. A., Zhukov, V., Tikhonovich, I., Tarakhovskaya, E., Frolov, A., & Wessjohann, L. A. (2018). Methodology of drought stress research: Experimental setup and physiological characterization. *International Journal of Molecular Sciences*, 19(12), 4089. <https://doi.org/10.3390/ijms19124089>

Rehman, A., Jingdong, L., Shahzad, B., Chandio, A. A., Hussain, I., Nabi, G., & Iqbal, M. S. (2015). Economic perspectives of major field crops of Pakistan: An empirical study. *Pacific Science Review B: Humanities and Social Sciences*, 1(3), 145–158. <https://doi.org/10.1016/j.psrb.2016.09.002>

Saha, D., Choyal, P., Mishra, U. N., Dey, P., Bose, B., Md, P., Gupta, N. K., Mehta, B. K., Kumar, P., Pandey, S., Chauhan, J., & Singh, R. K. (2022). Drought stress responses and inducing tolerance by seed priming approach in plants. *Plant Stress*, 4, 100066. <https://doi.org/10.1016/j.stress.2022.100066>

Sakamoto, A., & Murata, N. (2002). The role of glycine betaine in the protection of plants from stress: Clues from transgenic plants: Role of betaine in stress tolerance. *Plant, Cell & Environment*, 25(2), 163–171. <https://doi.org/10.1046/j.0016-8025.2001.00790.x>

Seleiman, M. F., Al-Suhaiibani, N., Ali, N., Akmal, M., Alotaibi, M., Refay, Y., Dindaroglu, T., Abdul-Wajid, H. H., & Battaglia, M. L. (2021). Drought stress impacts on plants and different approaches to alleviate its adverse effects. *Plants*, 10(2), 259. <https://doi.org/10.3390/plants10020259>

Shahbaz, M., Masood, Y., Perveen, S., & Ashraf, M. (2012). Is foliar-applied glycinebetaine effective in mitigating the adverse effects of drought stress on wheat (*Triticum aestivum* L.)?. *Journal of Applied Botany and Food Quality*, 84(2), 192.

Shanthi, N., Al-Huqail, A. A., Perveen, K., Vaidya, G., Bhaskar, K., Khan, F., & Alfaghām, A. (2023). Drought stress alleviation through nutrient management in *Cyamopsis tetragonoloba* L. *Journal of King Saud University - Science*, 102842. <https://doi.org/10.1016/j.jksus.2023.102842>

Tricker, P. J., ElHabti, A., Schmidt, J., & Fleury, D. (2018). The physiological and genetic basis of combined drought and heat tolerance in wheat. *Journal of Experimental Botany*, 69(13), 3195–3210. <https://doi.org/10.1093/jxb/ery081>


Wahab, A., Abdi, G., Saleem, M. H., Ali, B., Ullah, S., Shah, W., Mumtaz, S., Yasin, G., Muresan, C. C., & Marc, R. A. (2022). Plants' physio-biochemical and phyto-hormonal responses to alleviate the adverse effects of drought stress: A comprehensive review. *Plants*, 11(13), 1620. <https://doi.org/10.3390/plants11131620>

Weiskopf, S. R., Rubenstein, M. A., Crozier, L. G., Gaichas, S., Griffis, R., Halofsky, J. E., Hyde, K. J. W., Morelli, T. L., Morisette, J. T., Muñoz, R. C., Pershing, A. J., Peterson, D. L., Poudel, R., Staudinger, M. D., Sutton-Grier, A. E., Thompson, L., Vose, J., Weltzin, J. F., & Whyte, K. P. (2020). Climate change effects on biodiversity, ecosystems, ecosystem services, and natural resource management in the United States. *Science of The Total Environment*, 733, 137782. <https://doi.org/10.1016/j.scitotenv.2020.137782>

Yang, X., Lu, M., Wang, Y., Wang, Y., Liu, Z., & Chen, S. (2021). Response mechanism of plants to drought stress. *Horticulturae*, 7(3), 50. <https://doi.org/10.3390/horticulturae7030050>

Zhuang, J., Wang, Y., Chi, Y., Zhou, L., Chen, J., Zhou, W., Song, J., Zhao, N., & Ding, J. (2020). Drought stress strengthens the link between chlorophyll fluorescence parameters and photosynthetic traits. *PeerJ*, 8, e10046. <https://doi.org/10.7717/peerj.10046>

Zia, R., Nawaz, M. S., Siddique, M. J., Hakim, S., & Imran, A. (2021). Plant survival under drought stress: Implications, adaptive responses, and integrated rhizosphere management strategy for stress mitigation. *Microbiological Research*, 242, 126626. <https://doi.org/10.1016/j.micres.2020.126626>






Figure 1: Effect of *Glycine betaine* on (A) Shoot length, (B) shoot fresh weight under drought stress.







Figure 2: Effect of *Glycine betaine* on (A) Chlorophylla, (B) chlorophyll b and (c) carotenoids of wheat under drought stress.




Figure 3: Effect of *Glycine betaine* on (A) Photosynthetic rate (Pn), (B) transpiration rate (E), and (c) stomatal conductance (gs) of wheat under drought stress.

Submission: March 20, 2023, Revised: May 10, 2023

Accepted: May 30, 2023

Published Online: June 03, 2024